

# Sovaldi®

(sofosbuvir tablets)

## What is Sovaldi®?

- Sovaldi® is a prescription medication used for the treatment of hepatitis C virus.
- Sovaldi® is used in combination with other medications. Your doctor will inform you which medications you will take with Sovaldi.

## How should I take Sovaldi®?

- Take one tablet of Sovaldi® once daily, with or without food. **You should not take Sovaldi® alone.**
- Your doctor will tell you how long you should take Sovaldi® to cure your hepatitis C virus.

GENOTYPE 1 OR 4	12 WEEKS	SOVALDI, once a day	+	Ribavirin, twice a day	+	Peginterferon alfa, once a week
GENOTYPE 2	12 WEEKS	SOVALDI, once a day	+	Ribavirin, twice a day		
GENOTYPE 3	24 WEEKS	SOVALDI, once a day	+	Ribavirin, twice a day		

## What should I do if I miss a dose of Sovaldi®?

- If you forget to take a dose of Sovaldi®, take your dose as soon as you remember on the same day. **Do not take two tablets of Sovaldi® at the same time.** Take your next dose at your regular time the next day.

## Are there any side effects from Sovaldi®?

When taking Sovaldi® with ribavirin or Daklinza™, you may notice the following:

- ◆ Headache
- ◆ Tiredness

## What else should I know about Sovaldi®?

- **Do not start any new medications**, over-the-counter medications, vitamins, or herbal supplements without speaking to your healthcare team.
- Sovaldi® can **cause harm or death to your unborn baby** when taken with certain medications, such as ribavirin. You and your partner must use **2 forms of birth control during treatment and for at least 6 months after your treatment is complete. Do not become pregnant during this time.**
- It is important to have your blood tested during your treatment as instructed by your doctor.

