

Viekira Pak™

(ombitasvir, paritaprevir, ritonavir, and dasabuvir tablets)

What is Viekira Pak™?

Viekira Pak™ is a prescription medication used for the treatment of hepatitis C virus. Viekira Pak™ is a treatment which contains the following:

- Two combination tablets, which both contain ombitasvir, paritaprevir, and ritonavir. This is a pink tablet.
- Two dasabuvir tablets. This is a brown tablet.

How should I take Viekira Pak™?

- Viekira Pak™ is taken twice daily with food. You will take 3 tablets with your morning meal and 1 tablet with your evening meal. In some patients, you may need to take an additional tablet called ribavirin.
- Your doctor will tell you how long you should take Viekira Pak™ to cure your hepatitis C virus.



Are there any side effects from Viekira Pak™?

If you are taking Viekira Pak™ WITHOUT ribavirin, you may experience the following:

- ♦ Nausea
- ♦ Itching
- ♦ Difficulty sleeping

If you are taking Viekira Pak™ WITH ribavirin, you may experience the following:

- ♦ Nausea
- ♦ Itching
- ♦ Difficulty sleeping
- ♦ Tiredness
- ♦ Skin rashes/redness
- ♦ Feeling weak

If you notice yellowing of the eyes and/or skin, color changes in your stool, or if any of the above side effects get worse, call your doctor.

What else should I know about Viekira Pak™?

- **Do not start any new medications**, over-the-counter medications, vitamins, or herbal supplements without speaking to your healthcare team.
- Tell your doctor if you are using any medications containing ethinyl estradiol, such as birth control. You will need to use a different method of birth control during treatment and up to 2 weeks after your treatment is complete.
- It is important to have your blood tested during your treatment as instructed by your doctor.

What should I do if I miss a dose of Viekira Pak™?



Dasabuvir tablets:

Brown tablets, taken once in the morning and once in the evening, with food.

If you forget to take a dasabuvir tablet, take it as soon as you remember, **UNLESS** it has been **more than 6 hours** since your normal dose.

If it has been more than 6 hours, skip your missed dose and continue your treatment as normal.

Never take two dasabuvir tablets at once!

Ombitasvir/paritaprevir/ritonavir tablets:

Pink tablets. Two tablets are taken in the morning with food.

If you forget to take a ombitasvir / paritaprevir / ritonavir tablet, take it as soon as you remember, **UNLESS** it has been **more than 12 hours** since your normal dose.

If it has been more than 12 hours, skip your missed dose and continue your treatment as normal.